

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

Integral Field Spectroscopy of Extended Emission-Line Regions around QSOs

Hai Fu and Alan Stockton

*Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive,
Honolulu, HI 96822, USA*

Abstract. Luminous extended emission-line regions (EELRs) on kpc scales surround a substantial fraction of steep-spectrum radio-loud QSOs. Although their existence has been known for over three decades, there are still major uncertainties on the physical processes responsible for their complex morphology and kinematics. We are obtaining deep integral field spectroscopy for a sample of EELRs around QSOs at $z \leq 0.5$ with the Integral Field Unit (IFU) of the GMOS on the Gemini North telescope, aiming at extracting accurate kinematics of the EELRs, measuring important physical parameters (*e.g.*, density, temperature, metallicity) and reliable intensity ratios of diagnostic emission lines from individual clouds that comprise an EELR. Here we present results from the observations of the EELR of quasar 4C 37.43. We show maps of gas kinematics measured from the [O III] $\lambda 5007$ line and line-ratio diagnostic diagrams comparing the data with predictions from ionization models. We find that the ionized gas shows rather complex global kinematics, while linear velocity gradients are often seen in individual clouds. Pure photoionization by the QSO continuum is the most likely ionization mechanism for most of the EELR clouds.

About one third of low-redshift steep-spectrum radio-loud quasars are surrounded by massive ionized nebulae that often extend to several tens of kpc from the nucleus. These extended emission-line regions (EELRs) are important because they may reflect a key stage in a quasar's life, when it violently expels gas from its host galaxy (Stockton et al. 2002, 2006; Fu & Stockton 2006). Integral field spectrographs on large telescopes are powerful tools to study the EELRs. Here we present results from our recent study of the 4C 37.43 EELR, which is the most luminous EELR known among QSOs at $z \leq 0.5$.

4C 37.43 was observed with the GMOS/IFU in two configurations in the early half-nights of 2006 May 23 and 24 (UT): (1) Twelve exposures of 720 s were obtained with R831 grating and the full-field mode (IFU2) to cover $H\beta$ and [O III] $\lambda\lambda 4959, 5007$ lines. The telescope was dithered on a rectangular grid of $6'' \times 4''$ centered on the QSO, so the final FOV is $13'' \times 9''$. (2) Five 2400-s exposures were taken with B600 grating using the half-field mode (IFUR) to win a wider spectral coverage by losing half of the field (FOV shown in Fig. 1). The resulting datacube includes emission lines from [Ne V] $\lambda 3425$ to [O III] $\lambda 5007$.

The IFU datacubes (x,y, λ) carry a large amount of information. Due to the space limit, we discuss only the gas kinematics and the line-ratio diagnostics here. The full analysis will be presented elsewhere (Fu & Stockton, in prep.). (1) Gas kinematics were measured from the relatively isolated [O III] $\lambda 5007$ line in the mosaicked IFU2 datacube. As shown in Fig. 1, the EELR exhibits rather complex global kinematics. The relative line-of-sight velocities range from -700 to $+300$ km s $^{-1}$, and obviously most of the luminous clouds are blueshifted.




Figure 1. Velocity field of 4C 37.43 EELR derived from the [O III] $\lambda 5007$ line. When multiple velocities are present along the same line-of-sight, only the component with the lowest velocity is shown. Pixels are $0''.2$ squares. The crosses indicate the position of the quasar, which has been removed from the datacube using a PLUCY-based PSF subtraction technique. The dashed rectangle delineates the FOV of the IFUR datacube.




Figure 2. Line-ratio diagrams are overplotted with model grids. The pure photoionization model (Groves et al. 2004), shock-only and “shock + precursor” models (Dopita & Sutherland 1996) are plotted in black, blue and green, respectively. Measurements from various regions around 4C 37.43 are shown as blue circles, and red squares are data from four EELR clouds of 3C 249.1. Arrows are reddening vectors of $3 A_V$.

The velocity dispersions of most clouds are between 50 and 120 km s^{-1} . (2) The major ionization mechanism at work can be constrained by diagnostic diagrams involving ratios of some optical lines. Figure 2 shows examples of such diagrams. Line fluxes were measured from dereddened spectra (both Galactic and intrinsic reddenings have been taken into account), which were extracted either from the QSO-removed IFUR datacube or long slit spectra from Keck LRIS or DEIMOS. The associated $1\text{-}\sigma$ errors were obtained using a Monte-Carlo approach. These diagrams clearly show that the pure photoionization model best fits the data.

References

- Dopita, M. A., & Sutherland, R. S. 1996, *ApJS*, 102, 161
 Fu, H., & Stockton, A. 2006, *ApJ*, 650, 80
 Groves, B. A., Dopita, M. A., & Sutherland, R. S. 2004, *ApJS*, 153, 9
 Stockton, A., Fu, H., Henry, J. P., & Canalizo, G., 2006, *ApJ*, 638, 635
 Stockton, A., MacKenty, J. W., Hu, E. M., & Kim, T.-S. 2002, *ApJ*, 572, 735